



Inactivation Strategies for *Clostridium perfringens* Spores and Vegetative Cells

Prabhat K. Talukdar,^{a,b} Pathima Udombijitkul,^c Ashfaque Hossain,^d
Mahfuzur R. Sarker^{a,b}

Department of Biomedical Sciences, College of Veterinary Medicine, Oregon State University, Corvallis, Oregon, USA^a; Department of Microbiology, College of Science, Oregon State University, Corvallis, Oregon, USA^b; Department of Food Science and Technology, Faculty of Agro-Industry, Kasetsart University, Bangkok, Thailand^c; Department of Medical Microbiology and Immunology, Ras Al Khaimah Medical and Health Sciences University, Ras Al Khaimah, United Arab Emirates^d

ABSTRACT *Clostridium perfringens* is an important pathogen to human and animals and causes a wide array of diseases, including histotoxic and gastrointestinal illnesses. *C. perfringens* spores are crucial in terms of the pathogenicity of this bacterium because they can survive in a dormant state in the environment and return to being live bacteria when they come in contact with nutrients in food or the human body. Although the strategies to inactivate *C. perfringens* vegetative cells are effective, the inactivation of *C. perfringens* spores is still a great challenge. A number of studies have been conducted in the past decade or so toward developing efficient inactivation strategies for *C. perfringens* spores and vegetative cells, which include physical approaches and the use of chemical preservatives and naturally derived antimicrobial agents. In this review, different inactivation strategies applied to control *C. perfringens* cells and spores are summarized, and the potential limitations and challenges of these strategies are discussed.

KEYWORDS *Clostridium perfringens*, spores, vegetative cells, inactivation, antimicrobial agents, food poisoning

Clostridium perfringens is an anaerobic, spore-forming bacterium and can be found ubiquitously in the environment, including the guts of humans and other animals (1, 2). Although most strains of this species do not cause any harm to human and animals, a few of them are of concern due to their ability to cause a variety of histotoxic and gastrointestinal (GI) diseases (3, 4). *C. perfringens* can produce as many as 17 different toxins, but there is not a single strain that produces all of these toxins (3). Depending on the production of four major toxins (alpha, beta, epsilon, and iota), *C. perfringens* strains are classified into five toxin types (types A to E) (3). About 5% of global *C. perfringens* type A isolates produce *C. perfringens* enterotoxin (CPE), which is the major virulence factor for the pathogenesis of *C. perfringens*-associated food poisoning (FP) and nonfoodborne (NFB) GI diseases (3–6). Interestingly, most of the reported FP cases have been associated with *C. perfringens* isolates harboring the CPE-encoding gene (*cpe*) on the chromosome (C-*cpe* isolates), whereas isolates that possess *cpe* on the plasmid (P-*cpe* isolates) are linked to the occurrence of NFB GI diseases such as antibiotic-associated diarrhea and sporadic diarrhea, with some exceptions (7–9).

In addition to producing CPE, *C. perfringens* FP isolates have the ability to form spores that are highly resistant to various stress factors such as high temperature, high pressure, toxic chemicals, and radiation (10). However, *C. perfringens* vegetative cells are less resistant to these stress factors than their spore counterparts, and these resistance phenotypes vary among strains and growth conditions (11–13). For example, vegetative cells and spores of *C. perfringens* type A C-*cpe* isolates exhibited significantly higher

Accepted manuscript posted online 21
October 2016

Citation Talukdar PK, Udombijitkul P, Hossain A, Sarker MR. 2017. Inactivation strategies for *Clostridium perfringens* spores and vegetative cells. Appl Environ Microbiol 83:e02731-16. <https://doi.org/10.1128/AEM.02731-16>.

Editor Donald W. Schaffner, Rutgers, The State University of New Jersey

Copyright © 2016 American Society for Microbiology. All Rights Reserved.

Address correspondence to Mahfuzur R. Sarker, sarkerm@oregonstate.edu.

P.K.T. and P.U. contributed equally to this article.

heat resistance than those of P-*cpe* and *cpe*-negative isolates (11, 12, 14). Besides resistance to heat, vegetative cells and spores of C-*cpe* isolates possess higher resistance to various food preservative measures, including low temperature, osmosis-induced stress, and nitrite, than those of the P-*cpe* isolates (13, 15). These facts suggest the importance of developing an effective strategy that could kill or inhibit the growth of different strains of *C. perfringens* simultaneously.

The possession of high resistance properties facilitates the survival of *C. perfringens* spores in food vehicles, where they undergo germination and outgrowth to revert to vegetative cells and then cause FP illness in humans upon consumption of these contaminated foods (2). *C. perfringens* is currently ranked as the third most common foodborne pathogen in the United States (16), and it is estimated that almost a million cases of food-associated illnesses occurred annually in the United States, implicating *C. perfringens* as an etiological agent (17). This high incidence would subsequently lead to substantial losses in terms of economics, manpower, and medical expenses (18). Therefore, to minimize these substantial economical losses, strategies are being developed to inactivate the spores and vegetative cells of *C. perfringens*, especially in food industries and food products. In this review, we discuss the current status of different approaches to effectively inactivate the vegetative cells and spores of *C. perfringens*.

PHYSICAL APPROACHES

C. perfringens vegetative cells can easily be killed by introducing various abusive physical conditions, but their spore counterparts are difficult to inactivate with these treatments. However, studies have shown that either the manipulation of physical conditions or combined treatment with two or more stress factors was able to inactivate the spores (15, 19–21). Here, we discuss several of the physical parameters that can be used to inactivate spores and vegetative cells and their prospects for use in the food industry.

Thermal treatment. Thermal treatment is one of the most common ways of sterilizing the products, as excessive heat destroys the majority of the bacterial cells. It has been reported that *C. perfringens* spores are highly heat resistant (22), although the resistance patterns vary considerably with the strain and growth conditions, such as medium and incubation temperature (11). However, a significant inactivation of *C. perfringens* spores can be achieved by exposing them to high temperature for a longer period of time (23, 24). Wang et al. showed that more than 90% of *C. perfringens* spores were inactivated when they were incubated in water at 90 to 100°C for 10 to 30 min (23). Another report demonstrated that higher temperatures (110°C) were required to reduce the number of *C. perfringens* spores in the meat system (24) (Table 1). The introduction of other treatments (Table 1) in combination with moderate and high temperatures was also effective in the inactivation of both vegetative cells and spores of *C. perfringens* (19–21). For example, (i) implementation of ozone treatment followed by heat effectively inactivated both vegetative cells and spores of *C. perfringens* in a meat product (19), (ii) the simultaneous use of both thermal treatment and ultrasound or thermal treatment followed by ultrasound significantly increased the effectiveness of thermal inactivation of *C. perfringens* spores in a beef slurry (20), and (iii) pretreatment with gamma radiation followed by thermal treatment significantly decreased the number of spore counts (but interestingly, the reverse was not effective) (21).

Pressure treatment. High hydrostatic pressure (HHP) is one of the most widely used pasteurization techniques for food products other than temperature-dependent pasteurization. HHP treatment of food is more effective in killing of vegetative cells, and to some extent bacterial spores, than the conventional thermal processing of foods. It also retains the nutritional and sensory quality of the food products and consumes less energy than the other processing techniques (25). A combined treatment with HHP (650 MPa), temperature (75°C), and low pH (4.75) resulted in a 5.1-fold reduction of spores of *C. perfringens* type A P-*cpe* isolates but was not as effective (2.8-fold reduction) against spores of *C. perfringens* C-*cpe* isolates in laboratory medium (15). However, Gao et al. showed that the combined effects of HHP (654 MPa), heat (74°C), and nisin

TABLE 1 Effects of different inactivation strategies on the inhibition of *C. perfringens* cells and spores

Inactivation strategy	Matrix ^a	Treatment	Inhibitory effect	Reference(s)
Physical approaches				
Temp	Pork luncheon roll	1.3 min at 70°C for vegetative cells and 36 s at 110°C for spores	6-log reduction of spores and vegetative cells ^b	24
Temp and ozone	Beef surface	5 ppm aqueous ozone at 75°C	1–2-log CFU/g reduction of spores and vegetative cells	19
Temp and ultrasound	Beef slurry	Thermosonication (24 kHz, 0.33 W/g) at 75°C for 60 min	≤1.5-log reduction of spores	20
Temp and radiation	Buffer	Gamma radiation (0.7 Mrad) followed by thermal treatment at 103°C	6-log reduction of spores	21
HHP and temp	Poultry meat	586 MPa, 73°C for 10 min after spore germination with ≥50 mM L-asparagine and KCl mixtures	2–4-log CFU/g reduction of spore germination and outgrowth	103
HHP, pH, and temp	Buffer	650 MPa, 75°C at pH 4.75	1–5-fold reduction of spores for <i>C-cpe</i> isolates, 4–6-fold reduction for <i>P-cpe</i> isolates	15
Chemical agents				
Nitrate and nitrite	Broth	≥0.3% (wt/vol) NaNO ₂	1–4-log CFU/ml reduction in vegetative cells and spores	13
	Broth	100 ppm NaNO ₂ at 11°C	4-log CFU/ml reduction of spores and vegetative cells	30
	Ham	200 ppm NaNO ₂	2–4-log CFU/g reduction of spores and vegetative cells ^b	29
Sorbic acid	Buffer	1% (vol/vol) sodium sorbate	30–70% inhibition of spore germination for <i>C-cpe</i> isolates, and 50–90% inhibition for <i>P-cpe</i> isolates	39
Benzoic acid	Buffer	1% (vol/vol) sodium benzoate	30–70% inhibition of spore germination for <i>C-cpe</i> isolates, and 98% inhibition for <i>P-cpe</i> isolates	39
Lactic acid	Tajik-sambusa	2% (wt/wt) potassium lactate	1–2-log CFU/g reduction of spore germination and outgrowth ^b	44
	Injected pork	≥2% (wt/wt) calcium lactate, ≥3% (wt/wt) potassium lactate, or ≥3% (wt/wt) sodium lactate	4–6-log CFU/g reduction of spore germination and outgrowth ^b	45
	Injected turkey	≥2% (wt/wt) calcium lactate, ≥3% (wt/wt) potassium lactate, or ≥3% (wt/wt) sodium lactate	4–6-log CFU/g reduction of spore germination and outgrowth ^b	45
	Uncured ground turkey	2% (wt/wt) potassium lactate	2–4-log CFU/g reduction of spore germination and outgrowth ^b	42
	<i>Sous-vide</i> chicken	≥1.5% (wt/wt) sodium lactate	3–6-log CFU/g reduction of spore germination and outgrowth ^b	43
Acetic acid	Roast beef	≥2% MOstatin LV1	6–7-log CFU/g reduction of spore germination and outgrowth ^b	48
	Ground turkey roast	2.5% MOstatin V or 3.5% MOstatin LV	3–5-log CFU/g reduction of spore germination and outgrowth ^b	49
Phosphates	Poultry meat	1% (wt/vol) sodium tripolyphosphate	2–3-log CFU/g reduction of spore germination and outgrowth ^b	52

(Continued on next page)

TABLE 1 (Continued)

Inactivation strategy	Matrix ^a	Treatment	Inhibitory effect	Reference(s)
Natural antimicrobials				
Plant				
Cinnamaldehyde	Cooked ground beef and turkey	1% (wt/wt) cinnamaldehyde	3–4-log CFU/g reduction of spore germination and outgrowth ^b	61, 62
Carvacrol, thymol, and oregano oil	Cooked ground beef and turkey	2% (wt/wt) carvacrol, 2% (wt/wt) thymol, and 2% (wt/wt) oregano oil	3–5-log CFU/g reduction of spore germination and outgrowth ^b	61, 62
Tannins	Broth	≥5 mg/ml chestnut tannins	Up to 7-log CFU/ml reduction of vegetative growth	72
Green tea	Cooked ground chicken and pork	2% (wt/wt) green tea extracts	3–4-log CFU/g reduction of spore germination and outgrowth ^b	76
Animal				
Fatty acids	Buffer	1 mg/ml lauric acid	>7-log reduction of vegetative growth	79
Lysozyme	Broth	156 μg/ml lysozyme from hen egg white	Complete inhibition of vegetative cells	85
Chitosan	Cooked ground beef and turkey	3% (wt/wt) chitosan	4–5-log CFU/g reduction of spore germination and outgrowth ^b	90
Microbial				
Nisin	Broth	≥10 μM nisin	Complete inhibition of spore outgrowth and vegetative growth	92
Lacticin	Fresh pork sausage	2,500 AU/g lacticin 3147 with 2% (wt/vol) sodium lactate and/or 2% (wt/vol) sodium citrate	3–4-log CFU/g reduction of spore germination and outgrowth ^b	95

^aDifferent growth media were used for the inactivation strategies. Most of the experiments were performed with different meat products, as *C. perfringens* is a common pathogen in food products, especially in meats.

^bThe reduction value indicates the number of spores and/or cells compared to the total number of spores and/or cells of control samples (with no treatment).

(328 IU/ml) could effectively enhance the inactivation activity of HHP toward *C. perfringens* spores in milk products (26).

CHEMICAL AGENTS

Different types of chemical agents have been proven to be effective in controlling *C. perfringens*. Most of the studies have been focused on chemicals that can be used in food items as preservatives as well as flavor and color enhancers (27). Here, we discuss several of the common chemicals that have been used as food additives as well as food preservatives and have implications for the inhibition of *C. perfringens* spores and vegetative cells (Table 1).

Nitrate and nitrite. Nitrate and nitrite are well-known chemical preservatives for food to control spore-forming bacteria and have been used mostly in meat, fish, and cheese products for long periods of time (1, 27). These are also known as food additives, especially in cured meat, where nitrate and nitrite provide flavor and color stabilization (27). Nitrite can exert its antimicrobial effect against both vegetative cells and their spore counterparts, although spores are generally much more resistant to nitrite than growing cells (28). Nitrite affects the germination and outgrowth of *C. perfringens* spores in different types of meat products, and it has been reported that conventionally cured meat products, which have nitrite supplements, showed inhibition of the growth of *C. perfringens* (29, 30). However, commercially available organic meat products have the potential to support growth of *C. perfringens* (31, 32). Although nitrite is a very good antimicrobial agent, unfortunately its use in food products is limited due to its ability to form carcinogenic derivatives (33). This led to a search for alternative nitrite sources from natural compounds, although these compounds are not as effective as those used in conventionally cured meat for inhibition of the outgrowth of *C. perfringens* spores (34).

Organic acids. Organic acids are used as food preservatives and have been found to act as inhibitory agents for *C. perfringens* growth. Two of the most common and oldest organic acids used in food industries are sorbic acid (sorbate) and benzoic acid (benzoate). Both of these organic acids and their salts have been listed as generally recognized as safe (GRAS) compounds as food preservatives, with permissible concentrations approved by the U.S. Food and Drug Administration (FDA) (<http://www.accessdata.fda.gov/scripts/fdcc/?set=SCOGS>). Benzoates are one of the most widely used chemical antimicrobials because of their low cost, easy incorporation into foods, lack of impartment of color upon addition, and low toxicity (35, 36). However, sorbates are known mainly as antifungal agents. Tompkin et al. reported no difference in *C. perfringens* viability in presence or absence of potassium sorbate (37), but a follow-up study reported successful inhibition of *C. perfringens* growth by potassium sorbate in cooked sausage (38). Recently, our group showed the inhibitory effects of sorbate and benzoate derivatives against *C. perfringens* spore germination, outgrowth, and vegetative growth in both rich medium and a meat model system (39). Although the permissible levels of sorbate (0.3%, vol/vol) and benzoate (0.1%, vol/vol) could arrest *C. perfringens* spore outgrowth, higher concentrations (1%, vol/vol) are needed for the inhibition of spore germination in rich medium (Table 1) (39). Nevertheless, the permissible concentrations of sodium benzoate and potassium sorbate failed to control *C. perfringens* spore germination and outgrowth in cooked poultry meat during improper storage at 37°C, and this might result from the weak dissociation of the antimicrobials, rendering them ineffective against target microorganisms (39, 40).

Lactic acid and its derivatives of different salts have been reported to inhibit the germination and outgrowth of *C. perfringens* spores in different meat products, including injected turkey, injected pork, *sous-vide* chicken products, and tajik sambusa, under various abusive conditions (41–45). The lactate group represents a group of primary compounds responsible for the inhibition of germination and outgrowth of spores of *Clostridium* spp. (41) and could be utilized as an alternative to nitrite to warrant product safety during extended cooling of uncured meats (42). The addition of a 1.5% or higher concentrations of sodium lactate in marinated *sous-vide*-cooked chicken breast products led to the delay in the germination and outgrowth of spores of enterotoxigenic *C. perfringens* during storage at 19 and 25°C (43). Potassium lactate at 2% effectively restricted growth of a spore cocktail from 3 different *C. perfringens* strains during extended cooling of cooked uncured ground turkey breasts (42). However, calcium lactate was more effective than sodium lactate and potassium lactate in controlling germination and outgrowth of a spore cocktail from 3 *C. perfringens* strains in injected pork during a deviated chilling regimen (41).

Acetic acid, which is mostly used as vinegar, has been successfully applied for microbial growth inhibition and cell viability reduction for both Gram-positive and Gram-negative bacteria in different types of foods (46). Vinegar has been used as a condiment and food ingredient for the purpose of flavoring and preserving foods for thousands of years (47). A typical vinegar contains 5 to 40% acetic acid and other compounds that give the characteristic aroma (46). Recently, it was reported that a blend of vinegar and buffered lemon juice concentrate (MOStatin LV1) at different concentrations (0%, 2%, and 2.5%) was highly effective in controlling growth of *C. perfringens* spores in reduced-NaCl roast beef during abusive exponential cooling, regardless of the level of NaCl added and the cooling time used (48). The buffered vinegar (MOStatin V) at 2.5% and a blend of buffered lemon juice concentrate and vinegar (MOStatin LV) at 3.5% could also inhibit *C. perfringens* spore germination and outgrowth in ground turkey roast containing minimal ingredients (49).

Phosphates. Long-chain inorganic phosphates (polyphosphates [polyPs]) and the orthophosphate derivative blends have been extensively used in meat and dairy products for many years (50). These phosphates and their derivatives are used as food additives owing to their functional aspects regarding emulsification, stabilization, oxidation prevention, and flavor protection and, most importantly, as antimicrobial

TABLE 2 Antimicrobial activities of selected essential oils against *C. perfringens*

EO	MIC ($\mu\text{g/ml}$)	Reference
Cinnamaldehyde	140	60
<i>trans</i> -Cinnamaldehyde ^a	167	65
Thymol	375	63
Carvacrol	375	63
2- <i>tert</i> -Butyl-6-methyphenol ^a	175	65
Chamomile roman oil ^a	450	65
Citral oil ^a	275	65
Citronellal ^a	400	65
Geraniol ^a	450	65

^aThe MIC₉₅ is shown.

agents. Antimicrobial activities of variable-length phosphates (ortho-, pyro-, and polyphosphates) have been shown against many different bacteria, including *C. perfringens* (51–53). Our group evaluated the inhibitory effects of various polyPs on *C. perfringens* and demonstrated the following: (i) polyP significantly inhibited sporulation of *C. perfringens* by reducing sporulating cells by ~ 5 to $6 \log_{10}$; (ii) while *C. perfringens* spores were able to germinate in the presence of 1% sodium tripolyphosphate (STPP), their outgrowth was significantly inhibited; and (iii) STPP (at levels of $\geq 0.8\%$) exhibited antimicrobial activity against a spore cocktail from five different *C. perfringens* C-cpe isolates in chicken meat (52) (Table 1). Singh et al. found that the combination of sodium acid pyrophosphates from different sources were more effective than tetrasodium pyrophosphate in inhibition of *C. perfringens* growth in different meat products during abusive chilling conditions (53). However, the efficacy of these pyrophosphates depends on the type of meat product and dose of pyrophosphate (53).

NATURAL ANTIMICROBIALS

There are antimicrobial agents for inhibiting *C. perfringens* growth that are derived from natural sources, including animals, plants, and microorganisms (54, 55). Numerous researchers have studied the naturally occurring antimicrobial agents with regard to their properties of inhibition and inactivation of spoilage and disease-causing microorganisms (54, 55). The results revealed that many of them demonstrate a strong bactericidal effect and some of them are also inhibitory against spores of food-related bacteria; thus, there is a promising potential of these natural antimicrobial agents for being used as food preservatives. Table 1 listed some of the natural antimicrobials and the conditions for optimum inhibitory effects against *C. perfringens* spores and vegetative cells.

EOs. Essential oils (EOs) are secondary metabolites of aromatic plants having the characteristics of being volatile, natural, complex compounds with strong odors (56, 57). Historically, EOs have been used for thousands of years as food preservatives, for medicinal purposes, and for flavor and aroma. Many EOs have been shown to have antimicrobial activities against a broad range of bacteria, including *C. perfringens* (58). Cinnamaldehyde is one of the most common plant-derived EOs; it is listed as GRAS by the Flavor and Extract Manufacturers Association (FEMA) (59) and exhibited strong antibacterial activity against *C. perfringens* vegetative cells (Tables 1 and 2) (60). Cinnamaldehyde ($\geq 0.5\%$, wt/vol) was also effective in controlling *C. perfringens* spore germination and outgrowth in cooked ground beef and ground turkey during abusive chilling conditions (61, 62).

Carvacrol, thymol, and oregano oils are the key components in the EOs of oregano and thyme (58). The inhibitory effect of these EOs against germination and outgrowth of *C. perfringens* spores has been evaluated after deliberately contaminating cooked ground meats with *C. perfringens* spores prior to cooling in a deviated process from the U.S. Department of Agriculture Food Safety and Inspection Service (USDA-FSIS) stabilization guidelines for cooling meat products (61, 62). Carvacrol (1 to 2%, wt/vol), thymol (1 to 2%, wt/wt), and oregano oil (2%, wt/wt) effectively restricted *C. perfringens* growth from spores during 12 to 21 h of cooling from 54.4 to 7.2°C, with less than 1 log

CFU/g increase in the bacterial count in cooked ground beef (61). Another study, by Juneja and Friedman, showed that higher concentrations (2%, wt/wt) of carvacrol, thymol, and oregano oils were needed to completely restrict the growth of *C. perfringens* spores following more than 15 h of cooling of cooked ground turkey (62). This clearly indicates that the inhibitory effects of these EOs are variable and depend on the type and source of meat. Also, both thymol and carvacrol showed strong antibacterial activity against *C. perfringens* cells under *in vitro* conditions (Table 2) (63). Although *in vivo* studies with a chicken model showed that the supplementation of food with thymol and carvacrol did not reduce the number of *C. perfringens* in the chicken, it caused the reduction of the numbers of other bacteria and intestinal lesions in the gut and also enhanced the intestinal integrity and immune response (63, 64).

An extensive study by Si et al. with 66 different EOs demonstrated that 33 EOs inhibited >80% *C. perfringens* growth, while 9 EOs showed 50 to 80% inhibition and 24 EOs showed <50% inhibition (65). Out of the 33 EOs that exhibited >80% inhibition, seven were further investigated in a chicken model for their inhibitory potency against *C. perfringens* under acidic conditions. Excluding citral oil and geraniol, the inhibitory activities of the other five compounds (2-*tert*-butyl-6-methylphenol, carvacrol, chamomile roman oil, citronellal, and *trans*-cinnamaldehyde) were stable under acidic conditions (pH 2.0) (65). A few other studies also reported antimicrobial activity of some of the EOs against the *C. perfringens* growth (66–68). All these studies collectively suggest the potential use of EOs to reduce the number of *C. perfringens* organisms in food products.

Tannins. Tannins are water-soluble phenolic compounds that can be found in a variety of plants and are used as food additives or in leather processing industries due to their profound antimicrobial activity (69, 70). The effects of tannic acid and its hydrolytic product, gallic acid, were evaluated against vegetative cells of *C. perfringens* strain ATCC 13124 (71). The results showed that, tannic acid exhibited significant inhibition (lowest inhibitory concentration, 3.1 μ M) of *C. perfringens* growth at concentrations greater than 10 μ g/ml, while gallic acid showed no observable growth inhibition at concentrations up to 1,000 μ g/ml (71). This result suggests that the chemical structure of tannic acid is vital for retention of its antimicrobial activity. The source of tannins may also have an impact on their antimicrobial activity against *C. perfringens*. One study reported that the tannins from chestnut had higher antibacterial activity against *C. perfringens* than tannins from quebracho plants (72). However, tannins from both chestnut and quebracho could reduce the cytotoxicity of two *C. perfringens* toxins, alpha toxin and epsilon toxin, in MDCK cells (72). *C. perfringens* also showed no or very minimal resistance development against tannins over time compared to other antimicrobial products (73).

Green tea extracts. Green tea is one of the most consumed beverages worldwide and has beneficial effects on human health. It shows some antimicrobial activity against a wide range of bacteria, including *C. perfringens* (74). The activities of different polyphenols from extracts of green tea leaves against the growth of *C. perfringens* ATCC 13124 were evaluated by the disc diffusion method, and it was found that gallicocatechin and epigallocatechin showed the strongest growth inhibitory activity (75). The inhibitory effect of green tea extracts against germination and outgrowth of *C. perfringens* spores in different cooked meat products during deviated chilling regimens up to 21 h was also investigated (76). At 2.0%, the green tea extract with higher catechin content was successful in arresting growth from spores in all tested meats during a 21-h cooling period from 54.4 to 7.2°C (76). However, the efficacy of green tea extracts to prevent *C. perfringens* spore germination and outgrowth seems to be concentration and condition dependent (a higher dose of green tea extracts prevents *C. perfringens* growth in meat products for a longer storage time) (76). The catechin content of extracts is considered the key factor for the observed antibacterial activity, and this may partly explain the lack of activity of green tea extract with a lower total catechin content.

Fatty acids. Antimicrobial activity of short- and long-chain fatty acids against *C. perfringens* growth has been demonstrated in a few studies (77–79). However, long-chain fatty acids have been found to be more effective than short-chain fatty acids in inhibiting the growth of *C. perfringens* (77, 78). Saturated lauric acid (C_{12:0}) and unsaturated linoleic acid (C_{18:2}) at 0.5 mmol/liter were able to completely inhibit *C. perfringens* (77). A more recent study by Skøivanová et al. determined the susceptibility of vegetative cells of *C. perfringens* ATCC 13124 and CNCTC5459 (a clinical isolate) to different short- to long-chain fatty acids (79). Among them, caprylic (C_{8:0}), capric (C_{10:0}), lauric (C_{12:0}), myristic (C_{14:0}), and oleic (C_{18:0}) acids could effectively inhibit growth of both strains, and linoleic acid (C_{18:2}) inhibited only strain CNCTC5459 and not strain ATCC 13124 (79). Lauric acid (C_{12:0}) was proven to be the most potent antimicrobial, followed by myristic acid (C_{14:0}) and capric acid (C_{10:0}) (79). In an *in vivo* study, Timbermont et al. showed that the incidence of necrotic enteritis caused by *C. perfringens* in broiler chicken decreased if chickens were fed a diet supplemented with butyric acid in combination with lauric acid, thymol, cinnamaldehyde, and EO of eucalyptus at a lower dose (80).

Lysozyme. Lysozyme is a bacteriolytic enzyme that is obtained mainly from hen egg white and represents 3 to 5% of the egg albumin protein. It received GRAS status as a direct food additive and has been proven to be nontoxic to humans (36, 55). Lysozyme possesses a broad-spectrum antimicrobial activity and is commonly used in cheese production to prevent the late-blowing type of defect caused by *Clostridium* spp. (81, 82). Lysozyme inactivates bacterial cells by weakening the cell wall peptidoglycan (PG) layers via the hydrolysis of the β -1,4 linkages between *N*-acetylglucosamine and *N*-acetylmuramic acid, thus making it an effective antimicrobial agent against numerous Gram-positive bacteria (54, 83, 84). Lysozyme can inhibit the growth of *C. perfringens* vegetative cells at 156 μ g/ml (85) (Table 1). However, sublethal concentrations of lysozyme could inhibit the production of toxins (85). Based on the study of the effect of lysozyme on *Alicyclobacillus acidoterrestris*, it was hypothesized that lysozyme may inactivate spores by causing a rapid hydrolysis of the spore cortex, resulting in damage of the spore core (86). This notion may be supported by the fact that lysozyme can degrade the cortex of decoated *C. perfringens* spores lacking the cortex lytic enzyme SleC, leading to the resumption of wild-type-level colony-forming efficiency of *sleC* spores (87).

Chitosan. Chitosan is a biopolymer derived from the partial de-N-acetylation of chitin, which is the key component in the exoskeletons of crustaceans and is regarded as the second most abundant biopolymer in nature, following cellulose (88). It is nontoxic, nonantigenic, biocompatible, and biofunctional, and it has been approved as a food additive for many years. Moreover, its antimicrobial property makes it a potential food preservative in a variety of food products (89). A study by Juneja et al. demonstrated that 3% chitosan (degree of acetylation, 0.14) can reduce germination and outgrowth of spores of *C. perfringens* in cooked beef and turkeys by 4 to 5 log units during extended chilling regimens up to 18 h (90).

Nisin. Nisin, a class I bacteriocin and a polypeptide of 34 amino acid residues produced by certain strains of *Lactococcus lactis* subsp. *lactis*, has been successfully used as a food preservative in a variety of food products (91). It is approved by the FDA and WHO as GRAS and is permitted to be used in various food products in more than 50 countries. Nisin is an attractive option as a food additive because of its natural source and high effectiveness against a broad spectrum of Gram-positive bacteria. Several reports demonstrated the inhibitory activity of nisin alone or in conjunction with other food preservation technologies, such as heat and HHP, in controlling spores of *Clostridium* spp. in a variety of food products (26, 92). Nisin exerts its inhibitory effect against spore outgrowth and vegetative growth of *C. perfringens* FP and NFB isolates under laboratory conditions, but in a meat model system no such effect was observed (92). Nisin causes the lengthening of the lag phase of vegetative growth of *C. perfringens* type A strain NCTC8798 in a concentration-dependent manner (92). The higher the

concentration of nisin, the longer the lag phase observed. Nisin at 500 IU/ml or higher is thus required to prevent growth of *C. perfringens* NCTC8798 for up to 2 weeks (92).

Lacticin. Another type of class I bacteriocin is the two-component heat-stable lantibiotic lacticin 3147, produced by strains of *Lactococcus lactis* DPC 3147. Lacticin 3147 was reported to have a broad-spectrum inhibition toward Gram-positive bacteria, similar to that of nisin (93), but it is distinct from nisin in terms of its solubility and being active at physiological pH (94). It has been reported that the combination of lacticin and 2% sodium lactate was effective in reducing *C. perfringens* growth in fresh pork sausage (95).

Pediocin. Pediocins are well-known class II-type bacteriocins that are produced by *Pediococcus* spp. These are small, heat-stable, non-lanthionine-containing membrane active peptides (96). Pediocins are also affirmed as GRAS additives for application in certain food products (55). Different types of pediocins showed inhibitory activities against *C. perfringens* growth, such as pediocin AcH and pediocin PA-1 from *Pediococcus acidilactici* and pediocin A from *Pediococcus pentosaceus* (96–98). Pediocin PA-1 could lower the counts of *C. perfringens* by 2 and 0.8 log units, respectively, when frankfurters were stored at 10°C for 60 days and at 15°C for 30 days (97). Grilli et al. showed that a diet supplemented with pediocin A significantly improved the growth of broilers challenged with *C. perfringens* for up to 42 days (98).

GERMINATION-INDUCED INACTIVATION STRATEGIES

Having intrinsic resistance properties, *C. perfringens* spores can survive in the environment for extended periods (i.e., decades), and once these dormant spores come in contact with small molecules termed germinants, they return to being vegetative cells via the germination process (99, 100). In recent years, significant progress has been made in understanding the mechanism of *C. perfringens* spore germination, leading to the identification and characterization of suitable germinants and germinant receptors for spores of *C. perfringens* FP and NFB strains (99, 101). Although there are differences in the preference of germinants among the strains, some germinants (such as AK or L-cysteine) can induce germination of spores of a wide range of *C. perfringens* strains (99, 102). Such understanding has led to the development of novel strategies involving induction of spore germination followed by subsequent killing of germinated spores with mild treatments (39, 92, 103, 104), as in the following examples. (i) The application of germinant AK in meat products followed by HHP treatment (586 MPa) at high temperature (73°C for 10 min) significantly killed *C. perfringens* spores in the meat products (103) (Table 1). (ii) Permissive levels of chemical preservatives (such as nisin, sorbate, and benzoate) effectively arrested outgrowth of germinated *C. perfringens* spores in rich medium. However, higher levels of chemicals were needed to achieve significant inhibitory effects against *C. perfringens* spores inoculated into chicken meat (39, 91). (iii) Triggering spore germination considerably increased the sporicidal activity of commonly used disinfectants against *C. perfringens* FP spores attached to stainless steel chips (99). In addition, this germination-induced inactivation strategy also proved to be effective in killing spores from other *Clostridium* species (105–107). Collectively, triggering spore germination followed by inactivation treatment represents a novel strategy to enhance killing of *Clostridium* spores.

CHALLENGES OF INACTIVATION STRATEGIES AND FUTURE DIRECTIONS

Although there has been significant success in inactivating *C. perfringens* vegetative cells, the inactivation or elimination of *C. perfringens* spores from food products still remains a challenge. This is mostly because the effect of extreme physical stress damages the food quality and the use of excessive chemical preservatives results in toxicity and causes harm to human health. Furthermore, the addition of antimicrobial agents into food products might exhibit nonspecific inhibitory activity toward beneficial microorganisms harbored in the GI tract, causing an imbalance in the gut microflora (108, 109). These issues emphasize the importance of identifying the active compounds

and evaluating their toxicity prior to application to food products in order to warrant their safe usage. In addition, the elucidation of mechanisms of action of different physical and chemical techniques and naturally derived antimicrobials is critical for the efficient utilization of these treatments against the growth of *C. perfringens* spores and vegetative cells.

The recently developed germination-induced inactivation strategies also have some challenges (26, 103, 104). For example, due to spore germination heterogeneity, in a spore population some spores germinate within minutes and some takes hours or even longer to germinate (110). Therefore, further studies are needed to improve germination conditions for superdormant spores that exhibit slow or no germination and to validate the effectiveness of the germination-induced inactivation strategy under the practical conditions of food industry environments.

ACKNOWLEDGMENTS

This work was supported by grants from the N.L. Tartar Foundation and the Agricultural Research Foundation of Oregon State University (to M.R.S.).

REFERENCES

- Doyle MP, Buchanan RL (ed). 2013. Food microbiology: fundamentals and frontiers, 4th ed. ASM Press, Washington, DC. <https://doi.org/10.1128/9781555818463>.
- McClane BA, Uzal FA, Miyakawa MEF, Lyster D, Wilkins T. 2006. The enterotoxigenic clostridia, p 698–752. In Dworkin M, Falkow S, Rosenberg E, Schleifer K-H, Stackebrandt E (ed), The prokaryotes, 3rd ed, vol 4. Springer, New York, NY.
- Uzal FA, Freedman JC, Shrestha A, Theoret JR, Garcia J, Awad MM, Adams V, Moore RJ, Rood JI, McClane BA. 2014. Towards an understanding of the role of *Clostridium perfringens* toxins in human and animal disease. Future Microbiol 9:361–377. <https://doi.org/10.2217/fmb.13.168>.
- Brynstad S, Granum PE. 2002. *Clostridium perfringens* and foodborne infections. Int J Food Microbiol 74:195–202. [https://doi.org/10.1016/S0168-1605\(01\)00680-8](https://doi.org/10.1016/S0168-1605(01)00680-8).
- Xiao Y, Wagendorp A, Abee T, Wells-Bennik MH. 2015. Differential outgrowth potential of *Clostridium perfringens* food-borne isolates with various *cpe*-genotypes in vacuum-packed ground beef during storage at 12 °C. Int J Food Microbiol 194:40–45. <https://doi.org/10.1016/j.ijfoodmicro.2014.11.008>.
- Xiao Y, Wagendorp A, Moezelaar R, Abee T, Wells-Bennik MH. 2012. A wide variety of *Clostridium perfringens* type A food-borne isolates that carry a chromosomal *cpe* gene belong to one multilocus sequence typing cluster. Appl Environ Microbiol 78:7060–7068. <https://doi.org/10.1128/AEM.01486-12>.
- Collie RE, McClane BA. 1998. Evidence that the enterotoxin gene can be episomal in *Clostridium perfringens* isolates associated with non-food-borne human gastrointestinal diseases. J Clin Microbiol 36:30–36.
- Mpamugo O, Donovan T, Brett MM. 1995. Enterotoxigenic *Clostridium perfringens* as a cause of sporadic cases of diarrhoea. J Med Microbiol 43:442–445. <https://doi.org/10.1099/00222615-43-6-442>.
- Vaishnavi C, Kaur S. 2008. *Clostridium perfringens* enterotoxin in antibiotic-associated diarrhea. Indian J Pathol Microbiol 51:198–199. <https://doi.org/10.4103/0377-4929.41681>.
- Talukdar PK, Olguin-Araneda V, Alnoman M, Paredes-Sabja D, Sarker MR. 2015. Updates on the sporulation process in *Clostridium* species. Res Microbiol 166:225–235. <https://doi.org/10.1016/j.resmic.2014.12.001>.
- Sarker MR, Shivers RP, Sparks SG, Juneja VK, McClane BA. 2000. Comparative experiments to examine the effects of heating on vegetative cells and spores of *Clostridium perfringens* isolates carrying plasmid genes versus chromosomal enterotoxin genes. Appl Environ Microbiol 66:3234–3240. <https://doi.org/10.1128/AEM.66.8.3234-3240.2000>.
- Li J, McClane BA. 2006. Further comparison of temperature effects on growth and survival of *Clostridium perfringens* type A isolates carrying a chromosomal or plasmid-borne enterotoxin gene. Appl Environ Microbiol 72:4561–4568. <https://doi.org/10.1128/AEM.00177-06>.
- Li J, McClane BA. 2006. Comparative effects of osmotic, sodium nitrite-induced, and pH-induced stress on growth and survival of *Clostridium perfringens* type A isolates carrying chromosomal or plasmid-borne enterotoxin genes. Appl Environ Microbiol 72:7620–7625. <https://doi.org/10.1128/AEM.01911-06>.
- Novak JS, Juneja VK, McClane BA. 2003. An ultrastructural comparison of spores from various strains of *Clostridium perfringens* and correlations with heat resistance parameters. Int J Food Microbiol 86:239–247. [https://doi.org/10.1016/S0168-1605\(02\)00550-0](https://doi.org/10.1016/S0168-1605(02)00550-0).
- Paredes-Sabja D, Gonzalez M, Sarker MR, Torres JA. 2007. Combined effects of hydrostatic pressure, temperature, and pH on the inactivation of spores of *Clostridium perfringens* type A and *Clostridium sporogenes* in buffer solutions. J Food Sci 72:M202–M206. <https://doi.org/10.1111/j.1750-3841.2007.00423.x>.
- Lindstrom M, Heikinheimo A, Lahti P, Korkeala H. 2011. Novel insights into the epidemiology of *Clostridium perfringens* type A food poisoning. Food Microbiol 28:192–198. <https://doi.org/10.1016/j.fm.2010.03.020>.
- Scallan E, Hoekstra RM, Angulo FJ, Tauxe RV, Widdowson MA, Roy SL, Jones JL, Griffin PM. 2011. Foodborne illness acquired in the United States—major pathogens. Emerg Infect Dis 17:7–15. <https://doi.org/10.3201/eid1701.P11101>.
- Hoffmann S, Batz MB, Morris JG, Jr. 2012. Annual cost of illness and quality-adjusted life year losses in the United States due to 14 food-borne pathogens. J Food Prot 75:1292–1302. <https://doi.org/10.4315/0362-028X.JFP-11-417>.
- Novak JS, Yuan JT. 2004. Increased inactivation of ozone-treated *Clostridium perfringens* vegetative cells and spores on fabricated beef surfaces using mild heat. J Food Prot 67:342–346.
- Evelyn Silva FV. 2015. Use of power ultrasound to enhance the thermal inactivation of *Clostridium perfringens* spores in beef slurry. Int J Food Microbiol 206:17–23. <https://doi.org/10.1016/j.ijfoodmicro.2015.04.013>.
- Gombas DE, Gomez RF. 1978. Sensitization of *Clostridium perfringens* spores to heat by gamma radiation. Appl Environ Microbiol 36:403–407.
- Adams DM. 1973. Inactivation of *Clostridium perfringens* type A spores at ultrahigh temperatures. Appl Microbiol 26:282–287.
- Wang G, Paredes-Sabja D, Sarker MR, Green C, Setlow P, Li YQ. 2012. Effects of wet heat treatment on the germination of individual spores of *Clostridium perfringens*. J Appl Microbiol 113:824–836. <https://doi.org/10.1111/j.1365-2672.2012.05387.x>.
- Byrne B, Dunne G, Bolton DJ. 2006. Thermal inactivation of *Bacillus cereus* and *Clostridium perfringens* vegetative cells and spores in pork luncheon roll. Food Microbiol 23:803–808. <https://doi.org/10.1016/j.fm.2006.02.002>.
- Sarker MR, Akhtar S, Torres JA, Paredes-Sabja D. 2015. High hydrostatic pressure-induced inactivation of bacterial spores. Crit Rev Microbiol 41:18–26. <https://doi.org/10.3109/1040841X.2013.788475>.
- Gao Y, Qiu W, Wu D, Fu Q. 2011. Assessment of *Clostridium perfringens* spore response to high hydrostatic pressure and heat with nisin. Appl Biochem Biotechnol 164:1083–1095. <https://doi.org/10.1007/s12010-011-9196-0>.

27. Jay JM, Loessner MJ, Golden DA. 2005. Modern food microbiology, 7th ed, p 301–350. Springer, New York, NY.
28. Avila M, Gomez-Torres N, Hernandez M, Garde S. 2014. Inhibitory activity of reuterin, nisin, lysozyme and nitrite against vegetative cells and spores of dairy-related *Clostridium* species. *Int J Food Microbiol* 172:70–75. <https://doi.org/10.1016/j.jfoodmicro.2013.12.002>.
29. Redondo-Solano M, Valenzuela-Martinez C, Cassada DA, Snow DD, Juneja VK, Burson DE, Thippareddi H. 2013. Effect of meat ingredients (sodium nitrite and erythorbate) and processing (vacuum storage and packaging atmosphere) on germination and outgrowth of *Clostridium perfringens* spores in ham during abusive cooling. *Food Microbiol* 35:108–115. <https://doi.org/10.1016/j.fm.2013.02.008>.
30. Myers MI, Sebranek JG, Dickson JS, Shaw AM, Tarte R, Adams KR, Neibuh S. 2016. Implications of decreased nitrite concentrations on *Clostridium perfringens* outgrowth during cooling of ready-to-eat meats. *J Food Prot* 79:153–156. <https://doi.org/10.4315/0362-028X.JFP-15-301>.
31. Jackson AL, Sullivan GA, Kulchaiyawat C, Sebranek JG, Dickson JS. 2011. Survival and growth of *Clostridium perfringens* in commercial no-nitrate-or-nitrite-added (natural and organic) frankfurters, hams, and bacon. *J Food Prot* 74:410–416. <https://doi.org/10.4315/0362-028X.JFP-10-364>.
32. Jackson AL, Kulchaiyawat C, Sullivan GA, Sebranek JG, Dickson JS. 2011. Use of natural ingredients to control growth of *Clostridium perfringens* in naturally cured frankfurters and hams. *J Food Prot* 74:417–424. <https://doi.org/10.4315/0362-028X.JFP-10-365>.
33. Dahle HK. 1979. Nitrite as a food additive. *NIPH Ann* 2:17–24.
34. King AM, Glass KA, Milkowski AL, Sindelar JJ. 2015. Impact of clean-label antimicrobials and nitrite derived from natural sources on the outgrowth of *Clostridium perfringens* during cooling of deli-style turkey breast. *J Food Prot* 78:946–953. <https://doi.org/10.4315/0362-028X.JFP-14-503>.
35. Chipley JR. 2005. Sodium benzoate and benzoic acid, p 11–48. In Davidson PM, Sofos JN, Branan AL (ed), *Antimicrobials in food*, 3rd ed. Taylor & Francis Group, LLC, Boca Raton, FL.
36. Davidson PM, Taylor TM, Schmidt SE. 2013. Chemical preservatives and natural antimicrobial compounds, p 765–801. In Doyle MP, Buchanan RL (ed), *Food microbiology: fundamentals and frontiers*, 4th ed. ASM Press, Washington, DC. <https://doi.org/10.1128/9781555818463.ch30>.
37. Tompkin RB, Christiansen LN, Shaparis AB, Bolin H. 1974. Effect of potassium sorbate on *Salmonellae*, *Staphylococcus aureus*, *Clostridium perfringens*, and *Clostridium botulinum* in cooked, uncured sausage. *Appl Microbiol* 28:262–264.
38. Petaja E, Raevuori M, Poulanne E, Hill P. 1979. The effect of potassium sorbate and sodium nitrite on the organoleptic properties, stability, and growth of *Bacillus cereus* and *Clostridium perfringens* in cooked sausage. *Proc Eur Meet Meat Res Work* 25:917–921.
39. Alnoman M, Udombijitkul P, Paredes-Sabja D, Sarker MR. 2015. The inhibitory effects of sorbate and benzoate against *Clostridium perfringens* type A isolates. *Food Microbiol* 48:89–98. <https://doi.org/10.1016/j.fm.2014.12.007>.
40. Krebs HA, Wiggins D, Stubbs M, Sols A, Bedoya F. 1983. Studies on the mechanism of the antifungal action of benzoate. *Biochem J* 214:657–663.
41. Reddy Velugoti P, Rajagopal L, Juneja V, Thippareddi H. 2007. Use of calcium, potassium, and sodium lactates to control germination and outgrowth of *Clostridium perfringens* spores during chilling of injected pork. *Food Microbiol* 24:687–694. <https://doi.org/10.1016/j.fm.2007.04.004>.
42. Kennedy KM, Milkowski AL, Glass KA. 2013. Inhibition of *Clostridium perfringens* growth by potassium lactate during an extended cooling of cooked uncured ground turkey breasts. *J Food Prot* 76:1972–1976. <https://doi.org/10.4315/0362-028X.JFP-13-106>.
43. Juneja VK. 2006. Delayed *Clostridium perfringens* growth from a spore inocula by sodium lactate in *sous-vide* chicken products. *Food Microbiol* 23:105–111. <https://doi.org/10.1016/j.fm.2005.03.002>.
44. Yarbaveva SN, Velugoti PR, Thippareddi H, Albrecht JA. 2008. Evaluation of the microbial quality of Tajik sambusa and control of *Clostridium perfringens* germination and outgrowth by buffered sodium citrate and potassium lactate. *J Food Prot* 71:77–82.
45. Velugoti PR, Bohra LK, Juneja VK, Thippareddi H. 2007. Inhibition of germination and outgrowth of *Clostridium perfringens* spores by lactic acid salts during cooling of injected turkey. *J Food Prot* 70:923–929.
46. Ray B, Bhunia AK. 2008. Food biopreservatives of microbial origin, p 175–187. In Ray B, Bhunia AK (ed), *Fundamental food microbiology*, 4th ed. CRC Press, Boca Raton, FL.
47. Johnston CS, Gaas CA. 2006. Vinegar: medicinal uses and antiglycemic effect. *Med Gen Med* 8:61.
48. Li L, Valenzuela-Martinez C, Redondo M, Juneja VK, Burson DE, Thippareddi H. 2012. Inhibition of *Clostridium perfringens* spore germination and outgrowth by lemon juice and vinegar product in reduced NaCl roast beef. *J Food Sci* 77:M598–M603. <https://doi.org/10.1111/j.1750-3841.2012.02922.x>.
49. Valenzuela-Martinez C, Pena-Ramos A, Juneja VK, Korasapati NR, Burson DE, Thippareddi H. 2010. Inhibition of *Clostridium perfringens* spore germination and outgrowth by buffered vinegar and lemon juice concentrate during chilling of ground turkey roast containing minimal ingredients. *J Food Prot* 73:470–476.
50. Frenzel E, Letzel T, Scherer S, Ehling-Schulz M. 2011. Inhibition of cereulide toxin synthesis by emetic *Bacillus cereus* via long-chain polyphosphates. *Appl Environ Microbiol* 77:1475–1482. <https://doi.org/10.1128/AEM.02259-10>.
51. Obritsch JA, Ryu D, Lampila LE, Bullerman LB. 2008. Antibacterial effects of long-chain polyphosphates on selected spoilage and pathogenic bacteria. *J Food Prot* 71:1401–1405.
52. Akhtar S, Paredes-Sabja D, Sarker MR. 2008. Inhibitory effects of polyphosphates on *Clostridium perfringens* growth, sporulation and spore outgrowth. *Food Microbiol* 25:802–808. <https://doi.org/10.1016/j.fm.2008.04.006>.
53. Singh A, Korasapati NR, Juneja VK, Thippareddi H. 2010. Effect of phosphate and meat (pork) types on the germination and outgrowth of *Clostridium perfringens* spores during abusive chilling. *J Food Prot* 73:879–887. <https://doi.org/10.1016/j.jprot.2009.11.012>.
54. Davidson PM, Critzer FJ, Taylor TM. 2013. Naturally occurring antimicrobials for minimally processed foods. *Annu Rev Food Sci Technol* 4:163–190. <https://doi.org/10.1146/annurev-food-030212-182535>.
55. Juneja VK, Dwivedi HP, Yan X. 2012. Novel natural food antimicrobials. *Annu Rev Food Sci Technol* 3:381–403. <https://doi.org/10.1146/annurev-food-022811-101241>.
56. Bakkali F, Averbeck S, Averbeck D, Idaomar M. 2008. Biological effects of essential oils—a review. *Food Chem Toxicol* 46:446–475. <https://doi.org/10.1016/j.fct.2007.09.106>.
57. Guenther E, Althausen D. 1952. The essential oils, vol 6. Van Nostrand, New York, NY.
58. Burt S. 2004. Essential oils: their antibacterial properties and potential applications in foods—a review. *Int J Food Microbiol* 94:223–253. <https://doi.org/10.1016/j.jfoodmicro.2004.03.022>.
59. Adams TB, Cohen SM, Doull J, Feron VJ, Goodman JJ, Marnett LJ, Munro IC, Portoghese PS, Smith RL, Waddell WJ, Wagner BM. 2004. The FEMA GRAS assessment of cinnamyl derivatives used as flavor ingredients. *Food Chem Toxicol* 42:157–185. <https://doi.org/10.1016/j.fct.2003.08.021>.
60. Unlu M, Ergene E, Unlu GV, Zeytinoglu HS, Vural N. 2010. Composition, antimicrobial activity and *in vitro* cytotoxicity of essential oil from *Cinnamomum zeylanicum* Blume (*Lauraceae*). *Food Chem Toxicol* 48:3274–3280. <https://doi.org/10.1016/j.fct.2010.09.001>.
61. Juneja VK, Thippareddi H, Friedman M. 2006. Control of *Clostridium perfringens* in cooked ground beef by carvacrol, cinnamaldehyde, thymol, or oregano oil during chilling. *J Food Prot* 69:1546–1551.
62. Juneja VK, Friedman M. 2007. Carvacrol, cinnamaldehyde, oregano oil, and thymol inhibit *Clostridium perfringens* spore germination and outgrowth in ground turkey during chilling. *J Food Prot* 70:218–222.
63. Du E, Gan L, Li Z, Wang W, Liu D, Guo Y. 2015. *In vitro* antibacterial activity of thymol and carvacrol and their effects on broiler chickens challenged with *Clostridium perfringens*. *J Anim Sci Biotechnol* 6:58. <https://doi.org/10.1186/s40104-015-0055-7>.
64. Du E, Wang W, Gan L, Li Z, Guo S, Guo Y. 2016. Effects of thymol and carvacrol supplementation on intestinal integrity and immune responses of broiler chickens challenged with *Clostridium perfringens*. *J Anim Sci Biotechnol* 7:19. <https://doi.org/10.1186/s40104-016-0079-7>.
65. Si W, Ni X, Gong J, Yu H, Tsao R, Han Y, Chambers JR. 2009. Antimicrobial activity of essential oils and structurally related synthetic food additives towards *Clostridium perfringens*. *J Appl Microbiol* 106:213–220. <https://doi.org/10.1111/j.1365-2672.2008.03994.x>.
66. Lee JH, Cho S, Paik HD, Choi CW, Nam KT, Hwang SG, Kim SK. 2014. Investigation on antibacterial and antioxidant activities, phenolic and flavonoid contents of some thai edible plants as an alternative for

- antibiotics. *Asian-Australas J Anim Sci* 27:1461–1468. <https://doi.org/10.5713/ajas.2013.13629>.
67. Wannissorn B, Jarikasm S, Siriwanthai T, Thubthimthed S. 2005. Antibacterial properties of essential oils from Thai medicinal plants. *Fitoterapia* 76:233–236. <https://doi.org/10.1016/j.fitote.2004.12.009>.
 68. Kacaniová M, Vuković N, Horska E, Salamon I, Bobková A, Hleba L, Fiskelová M, Vatlak A, Petrova J, Bobko M. 2014. Antibacterial activity against *Clostridium* genus and antiradical activity of the essential oils from different origin. *J Environ Sci Health B* 49:505–512. <https://doi.org/10.1080/03601234.2014.896673>.
 69. Scalbert A. 1991. Antimicrobial properties of tannins. *Phytochemistry* 30:3875–3883. [https://doi.org/10.1016/0031-9422\(91\)83426-L](https://doi.org/10.1016/0031-9422(91)83426-L).
 70. Chung K-T, Murdock CA. 1991. Natural systems for preventing contamination and growth of microorganisms in foods. *Food Struct* 10:10.
 71. Chung KT, Lu Z, Chou MW. 1998. Mechanism of inhibition of tannic acid and related compounds on the growth of intestinal bacteria. *Food Chem Toxicol* 36:1053–1060. [https://doi.org/10.1016/S0278-6915\(98\)00086-6](https://doi.org/10.1016/S0278-6915(98)00086-6).
 72. Elizondo AM, Mercado EC, Rabinovitz BC, Fernandez-Miyakawa ME. 2010. Effect of tannins on the *in vitro* growth of *Clostridium perfringens*. *Vet Microbiol* 145:308–314. <https://doi.org/10.1016/j.vetmic.2010.04.003>.
 73. Redondo LM, Dominguez JE, Rabinovitz BC, Redondo EA, Fernandez Miyakawa ME. 2015. Hydrolyzable and condensed tannins resistance in *Clostridium perfringens*. *Anaerobe* 34:139–145. <https://doi.org/10.1016/j.anaerobe.2015.05.010>.
 74. Friedman M. 2007. Overview of antibacterial, antitoxin, antiviral, and antifungal activities of tea flavonoids and teas. *Mol Nutr Food Res* 51:116–134. <https://doi.org/10.1002/mnfr.200600173>.
 75. Ahn YJ, Kawamura T, Kim M, Yamamoto T, Mitsuoka T. 1991. Tea polyphenols—selective growth-inhibitors of *Clostridium* spp. *Agric Biol Chem* 55:1425–1426. <https://doi.org/10.1080/00021369.1991.10870770>.
 76. Juneja VK, Bari ML, Inatsu Y, Kawamoto S, Friedman M. 2007. Control of *Clostridium perfringens* spores by green tea leaf extracts during cooling of cooked ground beef, chicken, and pork. *J Food Prot* 70:1429–1433.
 77. Galbraith H, Miller TB, Paton AM, Thompson JK. 1971. Antibacterial activity of long chain fatty acids and the reversal with calcium, magnesium, ergocalciferol and cholesterol. *J Appl Bacteriol* 34:803–813. <https://doi.org/10.1111/j.1365-2672.1971.tb01019.x>.
 78. Sprong RC, Hulstein MF, Van der Meer R. 2001. Bactericidal activities of milk lipids. *Antimicrob Agents Chemother* 45:1298–1301. <https://doi.org/10.1128/AAC.45.4.1298-1301.2001>.
 79. Škořivanová E, Marounek M, Dlouhá G, Kaňka J. 2005. Susceptibility of *Clostridium perfringens* to C2–C18 fatty acids. *Lett Appl Microbiol* 41:77–81. <https://doi.org/10.1111/j.1472-765X.2005.01709.x>.
 80. Timbermont L, Lanckriet A, Dewulf J, Nollet N, Schwarzer K, Haesebrouck F, Ducatelle R, Van Immerseel F. 2010. Control of *Clostridium perfringens*-induced necrotic enteritis in broilers by target-released butyric acid, fatty acids and essential oils. *Avian Pathol* 39:117–121. <https://doi.org/10.1080/03079451003610586>.
 81. Gomez-Torres N, Avila M, Gaya P, Garde S. 2014. Prevention of late blowing defect by reuterin produced in cheese by a *Lactobacillus reuteri* adjunct. *Food Microbiol* 42:82–88. <https://doi.org/10.1016/j.fm.2014.02.018>.
 82. Wasserfall F, Teuber M. 1979. Action of egg white lysozyme on *Clostridium tyrobutyricum*. *Appl Environ Microbiol* 38:197–199.
 83. Chung W, Hancock REW. 2000. Action of lysozyme and nisin mixtures against lactic acid bacteria. *Int J Food Microbiol* 60:25–32. [https://doi.org/10.1016/S0168-1605\(00\)00330-5](https://doi.org/10.1016/S0168-1605(00)00330-5).
 84. Ross AI, Griffiths MW, Mittal GS, Deeth HC. 2003. Combining nonthermal technologies to control foodborne microorganisms. *Int J Food Microbiol* 89:125–138. [https://doi.org/10.1016/S0168-1605\(03\)00161-2](https://doi.org/10.1016/S0168-1605(03)00161-2).
 85. Zhang G, Darius S, Smith SR, Ritchie SJ. 2006. *In vitro* inhibitory effect of hen egg white lysozyme on *Clostridium perfringens* type A associated with broiler necrotic enteritis and its alpha-toxin production. *Lett Appl Microbiol* 42:138–143. <https://doi.org/10.1111/j.1472-765X.2005.01812.x>.
 86. Bevilacqua A, Corbo MR, Buonocore GG, Del Nobile MA, Sinigaglia M. 2007. Antimicrobial effectiveness of lysozyme against *Alicyclobacillus acidoterrestris*. *Adv Food Sci* 29:47–52.
 87. Paredes-Sabja D, Setlow P, Sarker MR. 2009. SleC is essential for cortex peptidoglycan hydrolysis during germination of spores of the pathogenic bacterium *Clostridium perfringens*. *J Bacteriol* 191:2711–2720. <https://doi.org/10.1128/JB.01832-08>.
 88. No HK, Meyers SP. 1995. Preparation and characterization of chitin and chitosan—a review. *J Aquat Food Prod Technol* 4:27–52. https://doi.org/10.1300/J030v04n02_03.
 89. No HK, Meyers SP, Prinyawiwatkul W, Xu Z. 2007. Applications of chitosan for improvement of quality and shelf life of foods: a review. *J Food Sci* 72:R87–R100. <https://doi.org/10.1111/j.1750-3841.2007.00383.x>.
 90. Juneja VK, Thippareddi H, Bari L, Inatsu Y, Kawamoto S, Friedman M. 2006. Chitosan protects cooked ground beef and turkey against *Clostridium perfringens* spores during chilling. *J Food Sci* 71:M236–M240. <https://doi.org/10.1111/j.1750-3841.2006.00109.x>.
 91. Delves-Broughton J. 2005. Nisin as a food preservative. *Food Aust* 57:525–527.
 92. Udompititkul P, Paredes-Sabja D, Sarker MR. 2012. Inhibitory effects of nisin against *Clostridium perfringens* food poisoning and nonfood-borne isolates. *J Food Sci* 77:M51–M56. <https://doi.org/10.1111/j.1750-3841.2011.02475.x>.
 93. Ryan MP, Rea MC, Hill C, Ross RP. 1996. An application in cheddar cheese manufacture for a strain of *Lactococcus lactis* producing a novel broad-spectrum bacteriocin, lacticin 3147. *Appl Environ Microbiol* 62:612–619.
 94. Rea MC, Clayton E, O'Connor PM, Shanahan F, Kiely B, Ross RP, Hill C. 2007. Antimicrobial activity of lacticin 3147 against clinical *Clostridium difficile* strains. *J Med Microbiol* 56:940–946. <https://doi.org/10.1099/jmm.0.47085-0>.
 95. Scannell AG, Ross RP, Hill C, Arendt EK. 2000. An effective lacticin biopreservative in fresh pork sausage. *J Food Prot* 63:370–375.
 96. Bhunia AK, Johnson MC, Ray B. 1988. Purification, characterization and antimicrobial spectrum of a bacteriocin produced by *Pediococcus acidilactici*. *J Appl Bacteriol* 65:261–268. <https://doi.org/10.1111/j.1365-2672.1988.tb01893.x>.
 97. Nieto-Lozano JC, Reguera-Useros JI, Peláez-Martínez MdC, Sacristán-Pérez-Minayo G, Gutiérrez-Fernández AJ, de la Torre AH. 2010. The effect of the pediocin PA-1 produced by *Pediococcus acidilactici* against *Listeria monocytogenes* and *Clostridium perfringens* in Spanish dry-fermented sausages and frankfurters. *Food Control* 21:679–685. <https://doi.org/10.1016/j.foodcont.2009.10.007>.
 98. Grilli E, Messina MR, Catelli E, Morlacchini M, Piva A. 2009. Pediocin A improves growth performance of broilers challenged with *Clostridium perfringens*. *Poult Sci* 88:2152–2158. <https://doi.org/10.3382/ps.2009-00160>.
 99. Paredes-Sabja D, Torres JA, Setlow P, Sarker MR. 2008. *Clostridium perfringens* spore germination: characterization of germinants and their receptors. *J Bacteriol* 190:1190–1201. <https://doi.org/10.1128/JB.01748-07>.
 100. Paredes-Sabja D, Setlow P, Sarker MR. 2011. Germination of spores of Bacillales and Clostridiales species: mechanisms and proteins involved. *Trends Microbiol* 19:85–94. <https://doi.org/10.1016/j.tim.2010.10.004>.
 101. Banawas S, Paredes-Sabja D, Korza G, Li Y, Hao B, Setlow P, Sarker MR. 2013. The *Clostridium perfringens* germinant receptor protein GerKC is located in the spore inner membrane and is crucial for spore germination. *J Bacteriol* 195:5084–5091. <https://doi.org/10.1128/JB.00901-13>.
 102. Udompititkul P, Alnoman M, Banawas S, Paredes-Sabja D, Sarker MR. 2014. New amino acid germinants for spores of the enterotoxigenic *Clostridium perfringens* type A isolates. *Food Microbiol* 44:24–33. <https://doi.org/10.1016/j.fm.2014.04.011>.
 103. Akhtar S, Paredes-Sabja D, Torres JA, Sarker MR. 2009. Strategy to inactivate *Clostridium perfringens* spores in meat products. *Food Microbiol* 26:272–277. <https://doi.org/10.1016/j.fm.2008.12.011>.
 104. Udompititkul P, Alnoman M, Paredes-Sabja D, Sarker MR. 2013. Inactivation strategy for *Clostridium perfringens* spores adhered to food contact surfaces. *Food Microbiol* 34:328–336. <https://doi.org/10.1016/j.fm.2013.01.003>.
 105. Nerandzic MM, Donskey CJ. 2010. Triggering germination represents a novel strategy to enhance killing of *Clostridium difficile* spores. *PLoS One* 5:e12285. <https://doi.org/10.1371/journal.pone.0012285>.
 106. Nerandzic MM, Donskey CJ. 2013. Activate to eradicate: inhibition of *Clostridium difficile* spore outgrowth by the synergistic effects of osmotic activation and nisin. *PLoS One* 8:e54740. <https://doi.org/10.1371/journal.pone.0054740>.
 107. Ishimori T, Takahashi K, Goto M, Nakagawa S, Kasai Y, Konagaya Y, Batori H, Kobayashi A, Urakami H. 2012. Synergistic effects of high

- hydrostatic pressure, mild heating, and amino acids on germination and inactivation of *Clostridium sporogenes* spores. *Appl Environ Microbiol* 78:8202–8207. <https://doi.org/10.1128/AEM.02007-12>.
108. Dorman HJ, Deans SG. 2000. Antimicrobial agents from plants: antibacterial activity of plant volatile oils. *J Appl Microbiol* 88:308–316. <https://doi.org/10.1046/j.1365-2672.2000.00969.x>.
109. Fabian D, Sabol M, Domaracka K, Bulnakova D. 2006. Essential oils—
- their antimicrobial activity against *Escherichia coli* and effect on intestinal cell viability. *Toxicol In Vitro* 20:1435–1445. <https://doi.org/10.1016/j.tiv.2006.06.012>.
110. Wang G, Zhang P, Paredes-Sabja D, Green C, Setlow P, Sarker MR, Li YQ. 2011. Analysis of the germination of individual *Clostridium perfringens* spores and its heterogeneity. *J Appl Microbiol* 111:1212–1223. <https://doi.org/10.1111/j.1365-2672.2011.05135.x>.

Prabhat K. Talukdar, M.S., is a Ph.D. student in the Department of Microbiology, Oregon State University, Corvallis, OR, USA. He obtained his B.Sc. in microbiology from the University of Dhaka (Dhaka, Bangladesh) in 2007, and in 2008 he obtained his M.S. in microbiology from the same university. He was a Research Officer at the Food and Enteric Microbiology Laboratory, International Center for Diarrheal Disease and Research, Bangladesh (ICDDR,B), from 2009 to 2011. He became interested in the bacterial genetics of enteropathogens during that period and started his doctoral thesis work at the Oregon State University in 2011. Currently his research focus is on the molecular mechanism of sporulation and germination of *Clostridium perfringens*.



Pathima Udombijitkul, Ph.D., received her Ph.D. in food science and technology from Oregon State University in 2013. She became a faculty member at the Department of Food Science and Technology, Faculty of Agro-Industry, Kasetsart University, Bangkok, Thailand, in 2013. Currently, she is a lecturer in various courses regarding food-associated microorganisms, including food microbiology, food safety, food plant sanitation, and hazard analysis and critical control points (HACCP). The spore-forming bacteria have attracted her interest for many years owing to their fascinating life cycle as well as their significance in human life, in both beneficial and harmful ways. She has been working on the basic biology of and inactivation strategies for sporeformers, particularly *Clostridium perfringens*, since 2007. Her current research areas involve the persistence of food-related sporeformers and the development of effective measures to control them.



Ashfaque Hossain, Ph.D., is a professor and chairperson of the Department of Medical Microbiology and Immunology at the Ras Al Khaimah Medical and Health Sciences University in Ras Al Khaimah, United Arab Emirates. He obtained his Ph.D. from the University of Glasgow, UK, and postdoctoral training at the University of Nebraska, Lincoln, NE, USA, and Harvard Medical School, Boston, MA, USA. His research interest is on antibiotic resistance mechanisms, biofilm production, and virulence mechanisms of enteric and foodborne pathogens, including *C. perfringens*.



Mahfuzur R. Sarker, Ph.D., is a professor in the Department of Biomedical Sciences, College of Veterinary Medicine, and Department of Microbiology, College of Science, Oregon State University, Corvallis, OR, USA. He obtained his Ph.D. from the University of Tokushima, Tokushima, Japan, and postdoctoral training from University Louvain, Brussels, Belgium, and the University of Pittsburgh School of Medicine, Pittsburgh, PA, USA. His research interests are (i) the molecular pathogenesis of the foodborne gastrointestinal pathogen *Clostridium perfringens*, (ii) mechanisms of sporulation/germination processes in *C. perfringens*, and (iii) the mechanism of *C. perfringens* spore inactivation.

